

**Sass-Klaassen, U. Leuschner, H.H., Buerkert, A. and Helle, G.**  
2008. Tree-ring analysis of *Juniperus excelsa* from the northern  
Oman mountains.

*In: Elferts, D., Brumelis, G., Gärtner, H., Helle, G. and Schleser, G.H.*  
(Eds.) *TRACE, Tree Rings in Archeology, Climatology and Ecology* 6,  
99 -108.

## Preface

This volume contains extended abstracts from talks and posters presented at the sixth TRACE (Tree Rings in Archaeology, Climatology and Ecology) conference, held in Riga (Latvia) May 3<sup>rd</sup> – 6<sup>th</sup>, 2007. The annual TRACE conference seeks to strengthen the network and scientific exchange between scientists and students involved in the study of tree rings. This annual conference is an initiative of the 'Association for Tree-Ring Research' (ATR). A high scientific level was maintained at the conference as at previous TRACE conferences, but simultaneously an informal forum was provided for young scientists and students to discuss concepts and ongoing or completed projects.

The conference was organized by the Latvian Dendroecologist Society and Faculty of Biology, University of Latvia. There were more than 80 participants at the conference from Australia, Austria, Belgium, Chile, Czech Republic, Estonia, Finland, France, Germany, Latvia, Lithuania, Poland, Romania, Slovenia, Spain, Sweden, Switzerland, and the United Kingdom. Outstanding oral and poster presentations were given by participants. In total, 36 talks were presented, covering the fields of Archaeology (5), Climatology (9), Ecology (7), Geomorphology (3), Isotopes (6), Methods (1), and Wood anatomy (5). In addition, 39 posters were displayed to the audience.

Two talks were given by invited speakers. Dr. Alar Läänelaid (University of Tartu, Estonia) presented an overview of dendrochronological studies in the three Baltic countries (Estonia, Latvia, Lithuania) to introduce the audience to the region, including ongoing and finished work as well as future perspectives for dendrochronological studies in the area. Dr. Hans Linderholm (Göteborg University, Sweden) presented a Summer North Atlantic Oscillation variability reconstruction over the last five centuries.

The editors of TRACE proceedings 2007 are delighted to present 25 short papers of tree-ring studies that were presented during the conference. We would like to thank the reviewers for their valuable comments on the first versions of these manuscripts. The organisers of the conference would like to thank sponsors (Administration of Latvian Environmental Protection Fund, State Stock Company „Latvia's State Forests”, Regent Instruments Inc.), whose contribution helped support the conference. Appreciation is also given to all participants for their valuable work, discussions, and exchange of ideas.

Didzis Elferts  
Guntis Brumelis  
Holger Gärtner  
Gerd Helle  
Gerhard Schleser



# CONTENTS

## SECTION 1 CLIMATOLOGY

<b><i>H.W. Linderholm, C.K. Folland &amp; J.W. Hurrell:</i></b>	8
Reconstructing Summer North Atlantic Oscillation (SNAO) variability over the last five centuries	
<b><i>J. Schultz, B. Neuwirth, D.A. Friedrichs, J. Löffler &amp; M. Winiger:</i></b>	17
Growth responses to NAO along a Central European west - east transect	
<b><i>S. Bogino &amp; F. Bravo:</i></b>	25
SOI and NAO impacts on <i>Pinus pinaster</i> Ait. growth in Spanish forests	
<b><i>A. Bräuning, P. von Schnakenburg, F. Volland-Voigt &amp; T. Peters:</i></b>	32
Seasonal growth dynamics and its climate forcing in a tropical mountain rain forest in southern Ecuador	
<b><i>D. Frank, O. Bouriaud, R. Wilson, G. Battipaglia, U. Büntgen, P. Fonti, K. Treydte, V. Trouet &amp; J. Esper:</i></b>	36
A challenge for spatially explicit reconstructions: the climate response of trees is a function of climate	
<b><i>U. Büntgen, D. Frank, K. Bareiss Valle, O. Bouriaud &amp; J. Esper:</i></b>	44
Climatic drivers of beech growth in the Vosges and Jura Mountains	
<b><i>J. Esper, R. Niederer, J. Luterbacher, U. Büntgen &amp; D. Frank:</i></b>	54
Calibration trails using very long instrumental and proxy data	
<b><i>V. Trouet &amp; J. Esper:</i></b>	61
Contrasting long-term drought signals in proxy records from northwestern Europe and the Mediterranean	
<b><i>Z. Fan &amp; A. Bräuning:</i></b>	68
Tree-ring based drought reconstruction in the central Hengduan Mountains region (China) since A.D. 1655	

## SECTION 2 DENDROECOLOGY

<b><i>D. Martín-Benito, P. Cherubini, M. del Río &amp; I. Cañellas:</i></b>	76
Differential effect of drought on <i>Pinus nigra</i> Arn. radial growth in mesic and xeric sites from southeastern Spain	

<b>J. Eckstein, H.H. Leuschner &amp; A. Bauerochse:</b>	83
Dendroecological studies on subfossil pine and oak from "Totes Moor" near Hannover (Lower Saxony, Germany)	
<b>Sz. Bijak:</b>	91
Various factors influencing the pointer year analysis	
<b>U. Sass-Klaassen, H.H. Leuschner, A. Buerkert &amp; G. Helle:</b>	99
Tree-ring analysis of <i>Juniperus excelsa</i> from the northern Oman mountains	

### SECTION 3 ISOTOPES

<b>J. Grießinger, A. Bräuning, A. Thomas &amp; G.H. Schleser:</b>	110
Stable oxygen isotopes in juniper trees from the Tibetan plateau as a proxy for monsoonal activity	
<b>P. von Schnakenburg, A. Bräuning &amp; G. Helle:</b>	115
Detecting annual growth rhythms from high-frequency densitometry and carbon isotopes in tropical mountain rain forest trees in southern Ecuador	
<b>O. Sidorova, T. Boettger, R. Siegwolf, M. Saurer, M. Naurzbaev &amp; E. Vaganov:</b>	121
Reaction of siberian subarctic larch trees to abrupt climatic changes derived from tree-ring and isotope data	

### SECTION 4 GEOMORPHOLOGY

<b>R.J. Kaczka, B. Wyzga &amp; J. Zawiejska:</b>	128
Tree-ring study of the island formation and riparian forest along a gravel-bed river in the Polish Carpathians	
<b>I. Malik &amp; D. Ciszewski:</b>	133
Meandering river bank erosion and channel lateral migration recorded in black alder ( <i>Alnus glutinosa</i> ) tree rings	

### SECTION 5 WOOD ANATOMY

<b>E. Přemyslovská, J. Šlezingerová &amp; L. Gandelová:</b>	140
Tree ring width and basic density of wood in different forest types	

## SECTION 6 ARCHAEOLOGY

- A. Vitas & M. Zunde:** 148  
Dendrochronological investigation on historical English oak (*Quercus robur* L.) in Lithuania and Latvia: problems and potential
- M. Rybníček, T. Kyncl, V. Gryc, E. Přemyslovská & H. Vavrčík:** 152  
Building of the oak standard chronology for the Czech Republic
- M. Opała & R.J. Kaczka:** 160  
Dating of wooden shelters in Polish High Tatras - tree rings records of the shepherding history in Carpathians

## SECTION 7 NEW APPLICATIONS

- J. König, B. Günther & C.T. Bues:** 168  
Application of multivariate cross-dating to historical timbers with less than 50 tree-rings from the Albrechtsburg Castle and the Meissen Cathedral, Saxony
- H. Vavrčík & V. Gryc & M. Rybníček:** 176  
Fluorescence microscopy utilization for lignin detection in wooden cell walls in spruce. A technical note
- W. Beck:** 183  
Mean growth behaviour of forest stands – Methodological aspects from dendrochronology and forest mensuration
- List of participants** 192  
TRACE 2007 Conference, May 3<sup>rd</sup> – 6<sup>th</sup> 2007, organized by the Latvian Dendroecologist Society and Faculty of Biology, University of Latvia, Riga, Latvia.



# Tree-ring analysis of *Juniperus excelsa* from the northern Oman mountains

U. Sass-Klaassen<sup>1</sup>, H.H. Leuschner<sup>2</sup>, A. Buerkert<sup>3</sup> & G. Helle<sup>4</sup>

<sup>1</sup> Forest Ecology and Forest Management (FEM), Wageningen University and Research Centre, P.O. Box 47, 6700 AA Wageningen, The Netherlands

<sup>2</sup> Laboratory for Dendroecology, Department of Palynology and Climate Dynamics, Albrecht-von-Haller Institute of Plant Science, University of Goettingen, Wilhelm-Weber-Str. 2a, 37073 Goettingen, Germany

<sup>3</sup> Organic Plant Production and Agroecosystems Research in the Tropics and Subtropics, University of Kassel, 37213 Witzenhausen, Germany

<sup>4</sup> Forschungszentrum Jülich GmbH, ICG-5, 52425 Jülich, Germany  
Email: Ute.SassKlaassen@wur.nl

## Introduction

### *Background*

Oman has a rich history of irrigation agriculture which was largely shaped by changes in climate conditions throughout the last millennia. The historical development of oasis agriculture in the mountainous north of the country reflects changes in settlement patterns that are thought to be strongly related to recurrent droughts (Siebert et al. 2004). To study (i) the driving role of climate, i.e. rainfall for the oasis agriculture in the past and (ii) the effects of possible changes in climate on the still existing oases long climate records are needed. However, for the whole Arabian Peninsula such records are scarce and – if available - only date back until the last century, when the British established the first weather stations in the region.

The only published climate archive which fulfils the requirements of longevity and high resolution originates from a varved sediment core of the NE Arabian Sea (von Rad et al. 1999). This 5,000-yr record was dated at an extraordinary precision (about one order of magnitude smaller than typical AMS <sup>14</sup>C data) and allowed to reconstruct the strength of both the winter and the late summer monsoon affecting the area. Nevertheless this record may be of only limited value for Oman with its more arid climate largely influenced by the Indian Summer Monsoon. The recent stalagmite records of Fleitmann et al. (2007) in contrast are from northern Oman and cover a timespan of 10,000 years but their resolution is limited.

### *Potential of Dendrochronology*

Dendrochronological studies on long-live tree species have the potential to provide long time series that reflect changes in climate conditions. In the high altitude Al Jabal al Akhdar mountains of Oman extended *Juniperus excelsa* stands exist growing under suitable site conditions. Tree-ring analysis of comparable material has been successfully conducted in both tropical and dry (mountainous) forests in Africa (Fichtler et al. 2004; Couralet et al. 2005; Schöngart et al. 2006, Trouet et al. 2006). Based on initial observations we assumed that these juniper trees form annual rings as they grow under harsh climatic conditions in the dry mountainous forests with a single pronounced dry season. Also a preliminary study by

Fisher et al. (1994) on *J. excelsa* from Oman suggested that *Juniperus excelsa* has potential for dendrochronology.

However, uncertainty remained about the periodicity of ring formation in trees growing under extremely dry conditions (Wils & Eshetu 2007) Theoretically both false or double rings and missing rings may occur in juniper (Couralet et al. 2005). False rings may be caused by a bimodal rainfall distribution in certain years due to winter rains and a strong monsoon event in any one year. Missing or partly missing rings (=wedging rings), in contrast, are frequently formed during years with little rainfall.

### Objectives

In this study we therefore addressed the following research questions: (1) Does *Juniperus excelsa* from Oman form clearly distinguishable growth layers? (2) What is the mean annual growth level and estimated age of juniper in the study area? (3) Is cross-dating of tree-ring series between different trees possible? And ultimately (4) what is the potential of *Juniperus excelsa* for the reconstruction of climate records?

## Materials and methods

### Study site and sample material

The sampling area is located in the Jabal al Akhdar mountain range in the northern part of Oman (Fig. 1a). Unfortunately, the only climate records from Oman are available from the coast at Muscat (Fig. 1b). The climate diagram from Muscat indicates a dry seasonal climate with a prolonged dry season from April to mid November during which drought conditions can be expected.




Figure 1: Location with the old *Juniperus excelsa* stands (see marked square on the insert) on Al-Jabal-al-Akhdar mountain range of northern Oman (left). Climate diagram from Muscat (Seeb Airport, right).

The climate at the high mountain study location is considerably cooler with a long term annual average air temperature of 18.1°C, a minimum of -3.6°C and a maximum of 36.3°C, the coolest and hottest month being January and June, respectively (Department of Civil

Aviation and Meteorology, Muscat, Oman). The juniper trees (*Juniperus excelsa* subsp. *Polycarpus*) are growing at about 2,300 m a.s.l. in an open dry mountainous forest. The trees are strongly protected which makes cutting of trees for sample collection impossible. However, in addition to 12 cored trees, from which two cores were taken at breast height in January 2004, four stem disks were available from trees cut due to recent road construction.

#### *Cross-dating of tree-ring series*

According to the above mentioned tree ring uncertainties (double rings, missing rings), the measured tree-ring series were visually crossdated in a stepwise hierarchical order: Starting with comparisons of series from the cores of the same tree we identified disagreements and went back to the wood surface to check the wood structure in the problematic area. In order to adjust the series, the single tree-ring series were edited in TSAP by adding missing rings and merging false or double rings. The corrected tree-ring series were then averaged into *tree chronologies*, which were visually checked and corrected as in the previous tree-level-check. After successful cross-dating the corrected tree-ring series were combined into a *site chronology*. The final result of the visual cross-dating were statistically verified by running correlation analysis performed with the software program COFECHA (Holmes 1983).

#### *<sup>14</sup>C dating*

As tree-ring formation in juniper from Oman is expected to be very spurious we decided to use <sup>14</sup>C dating to check whether dendrochronological detection of tree rings coincides with the age suggested by <sup>14</sup>C dating. For that purpose <sup>14</sup>C dating was done on the 10 innermost tree rings of the three samples with most tree rings (111, 105, disk 27 in table 1) and compared to the result of the dendro-dating after measuring and cross dating of two radii.

#### *Age determination and statistical description of tree-ring series*

The number of measured (and corrected) tree rings indicates the minimum tree age. An exact age determination is, however, only possible if cores include the pith (=centre of the tree) and if reliable corrections for the missing rings according to the sampling heights can be made. The mean tree-ring width (and the standard deviation) describes the general growth level and the variability in tree-ring width and gives an indication about the chance of missing rings in the tree-ring record. Mean sensitivity and autocorrelation of first order are standard parameters in dendrochronology to describe the changes in tree-ring width from year to year and the persistence of the tree-ring pattern (Cook & Kairiukstis 1990).

#### *$\delta^{18}\text{O}$ analyses*

Many tree-ring series show alternating phases of high-frequency dominated or low-frequency dominated variation with the enhanced high-frequency signal being caused by the regular occurrence of very narrow rings. It is, however, well possible that these narrow rings resemble density variations, that is second growth phases that occur in periods with a strong bimodal rainfall distribution (Fig. 1b). This growth behaviour has been proved for many species growing under dry condition, such as in the Mediterranean's where temporary dry

spells cause a bimodal growth pattern with a recovery of growth after a summer drought (Cherubini et al. 2003). If so, changes in spectral behaviour of the tree-ring series can be used to point out shifts in large-scale climate-driving factors such as monsoon activity that trigger the amount and the distribution of rainfall throughout the year.  $\delta^{18}\text{O}$  is known to be a sensitive indicator of changes in water availability and hence rainfall patterns. To study whether or not changes in the spectral behaviour of a tree-ring series are corresponding with the isotopic ( $\delta^{18}\text{O}$ ) composition,  $\delta^{18}\text{O}$  was measured in altogether 50 ring layers of sample 27. 25 consecutive tree rings bridging 1905 to 1930 the change from a high-frequency signal period to a period with moderate variability and b) 25 layers from the more recent (moderate as well) period 1975 to 2000.

## Results and Discussion

*Does Juniperus excelsa from Oman form clearly distinguishable growth layers?*

*Wood anatomical characteristics of Juniperus excelsa*

As in case of conifers from regions with non-ambiguous annual tree rings, the growth rings in juniper are mostly defined by the sharply edged contrast between the last few rows of flattened and thick-walled late-wood tracheids and the larger and thin-walled early-wood tracheids of the next growth ring. These rings could be clearly identified and measured. However, some small rings consisted of only a few cell rows of early and late-wood with a less typically contrast in cell size but clear and sharp edges to the previous or next ring. These layers could not be clearly identified as density fluctuations, false rings or real “annual” rings (Fig. 2).



*Figure 2: Cross section of Juniperus excelsa from with distinct tree rings. The dark part (late wood) and light part (early wood) constitute a single annual growth layer (left). Partly missing rings and density variations in Juniperus excelsa (right).*

*What is the mean annual growth level and estimated age of juniper in the study area?*

The growth of the juniper trees from Al Jabal al Akhdar was very slow with an average of 0.52 mm (Tab. 1). However, it varied considerably between trees. The high variability indicates that growth conditions of the juniper trees are affected by different micro-site conditions such as surface slope-dependent water supply, matching the observations by Fisher & Gardner (1995).

**Table 1.** (Statistical) description of tree-ring series. No. rings = number of measured tree rings, Mean TRW = mean tree-ring width [mm], Stdv = Standard deviation, AC(1)= 1<sup>st</sup> order autocorrelation, MS = mean sensitivity.

Tree no.	Sampling date	No. rings	Mean TRW	Stdv	AC(1)	MS(%)
101	2004	214	1.37	1.154	0.02	91
102	2004	207	0.57	0.387	0.35	64
103	2004	172	0.58	0.357	0.43	51
104	2004	364	0.52	0.334	0.10	69
105	2004	604	0.27	0.144	0.21	55
106	2004	306	0.57	0.405	0.10	72
107	2004	392	0.34	0.173	0.20	53
108	2004	213	0.27	0.163	0.40	51
109	2004	482	0.24	0.141	0.52	45
110	2004	387	0.67	0.508	0.13	82
111	2004	783	0.21	0.154	0.34	59
112	2004	267	0.38	0.229	0.20	65
Disk 1	2005	306	0.46	0.415	0.34	64
Disk 2	2005	100	1.10	1.132	0.49	56
Disk 3	2005	146	0.48	0.303	0.27	59
Disk 27	2003	622	0.39	0.301	0.23	71

The number of measured tree rings provided an indication of minimum tree (pith-) age and varied considerably between the sample trees with tree 111 being the oldest tree (783 rings) with the lowest growth level and tree 'disk 2' being the youngest tree with 100 measured rings. However, as most samples did not include the pith, an unknown number of inner rings might have been omitted. The number of measured tree rings shows that junipers in this area can get very old which makes them highly valuable for climate reconstruction. Nevertheless, it has to be noticed that radial growth – especially in old individuals - under these extremely harsh climate and edaphic conditions can be extremely slow and might even stop for several (consecutive) years.

#### *Comparison between <sup>14</sup>C dating and tree ring measurements*


The <sup>14</sup>C dating of the juniper sample showed that there was at least – according to the youngest border of the 95.4 % probability level, Oxcal Calibration - a difference between the <sup>14</sup>C date and the dendro date of 30, 25 and 20 tree rings, respectively, for the three samples covering 783, 622 and 604 years. This indicates that some more missing rings have to be expected in the measured tree-ring series.

### *Is cross-dating of tree-ring series between different trees possible?*

#### *Tree-ring characteristics and tree-ring patterns*

Tree-ring formation in the juniper trees from Al Jabal al Akhdar was very irregular with abrupt changes between wider rings and extremely small rings (Fig. 2). Wedging, that is partly missing rings, was frequently visible on the cores which suggests that a considerable number of missing rings can be expected (Fig. 2). Double rings, that is density variations, only occurred on wide tree rings (>1.5 mm). This leads to a generally high sensitivity and low autocorrelation in tree-ring series (Tab. 1).

Unfortunately the tree-ring patterns from two cores of the same tree differed considerably. This was likely due to the eccentric growth pattern of the trees whereby wood formation around the circumference was very variable and made tree-ring analysis based on two cores difficult (Fig. 2). Cross-dating between two (or three) cores of the same tree showed that sometimes up to 30 rings were missing in one tree-ring record. Missing rings mainly occur during phases of depressed growth. However, we also observed that tree rings disappeared during phases of apparently normal growth. Given the fact that double rings most often occurred in wide tree rings, it was assumed that shifts in tree-ring series are mainly due to missing rings. The tree-ring series were corrected by tracing well-matching parts in two tree-ring series and fitting tree rings in until both series matched well. This extremely time-consuming process was applied to all 12 samples and the samples of the stem disks separately. Eventually all samples were matched and combined into a site chronology.



*Figure 3: Cross-dated tree-ring series of 12 juniper trees from Al-Jabal-al-Akhdar together with the chronology (shifted, on top).*

#### *Calculation of a (preliminary) site chronology and description of tree-ring series*


Figure 3 shows the cross-dated and corrected tree-ring series of 12 juniper trees (trees101-112) with the site chronology on top. The fact that such a site chronology could be developed

demonstrates that the tree-ring series of single trees show periods of striking similarity. Figures 3 illustrates the tree-ring patterns in the period from 1800 to 2003 only. Cross-dating was possible throughout the whole period where at least two tree-ring series overlapped (1400-2004).

The single tree-ring series exhibited no clear age trend. Instead all trees showed a very sensitive growth pattern whereby phases of wider tree rings alternated with depressed tree growth. Obvious growth depressions occurred in the 1860ies, around 1915 and around 1980. However, the trees were apparently able to recover from these depressions and there also was no indication of a prolonged decline in growth activity in recent decades.

#### *Changes in spectral behaviour – $\delta^{18}\text{O}$ isotope composition*

In single tree ring series striking changes in spectral behaviour of tree-ring widths could be detected. Phases of high sensitivity alternate with phases of dominant low frequency. In can be seen in sample 27 (Fig. 4), where a slight tendency towards more decadal variations was observed after the major depression in c. 1910. However, also in the chronology it was possible to distinguish phases that either contained more high frequency (annual) variation or more decadal variation (Fig. 3). In view of this, the  $\delta^{18}\text{O}$  analyses on two 25-year periods in sample 27 yielded interesting results. The major switch in spectral behaviour of the tree-ring width, recorded around 1913, was related to a shift in the  $\delta^{18}\text{O}$ -isotope composition towards significantly higher values known to reflect more humid conditions. The isotope composition of the second sample taken in the more recent period from 1975 to 2000 shows the same high values of  $\delta^{18}\text{O}$ . Although these results need much more support by additional studies they indicate that frequency changes in tree-ring width are related to changes in the hydrological regime, which in turn could be related to changes in the amount or distribution of precipitation throughout the year.



*Figure 4. Typical growth pattern with small rings in 2-3 layers distance during a high frequency growth period. The interpretation of the small layers as annual rings, double rings or extreme density fluctuations is unclear (left). Tree-ring series of sample 27 with a shift between a phase with high frequency variation due to occurrence of small rings and more low- frequency variation. The  $\delta^{18}\text{O}$  composition data (above curves) of two 25-year segments indicate a strong and persistent increase in humidity from c. 1915 onwards (right).*

### *What is the potential to use Juniperus excelsa for climate reconstruction?*

The calculated site chronology has to be regarded as preliminary. It certainly needs further support by integrating more material from – preferably - stem disks. However, as the juniper trees are strongly protected in Oman it is not likely that it will be possible to collect enough such material to considerably complement our samples. But even if stem-disk collection would not be such a major problem, it is still questionable if this material is appropriate to build up long tree-ring chronologies with the classical dendrochronological techniques that have been used in these and other studies. Juniper from Oman has proved to be very difficult material for several reasons: (1) the trees are generally slow growing; (2) show a very irregular growth pattern with alternating phases of sensitive (mainly annual variation) and very complacent (mainly decadal variation) growth; real tree-ring boundaries are often difficult to distinguish from density variations, that is double rings frequently occur; (4) (partly) missing rings are a very common phenomenon. The last point causes major trouble and the  $^{14}\text{C}$  analyses suggested that even after thorough crossdating, that already resulted in tracing many (partly) missing rings, there still were rings that could not be detected.

Taking into account these aspects the following recommendations can be made:

Tree-ring analysis with juniper from Oman needs support by additional techniques.  $^{14}\text{C}$  Wiggle matching (Kuzmin et al. 2004) is essential in order to check whether all missing trees can be traced via cross dating of tree-ring series. A carefully selection of samples, preferably stem disks, that undergo extensive wiggle matching could provide the exact time frame for verifying dendrochronological dating.

Once the preliminary chronology is supported by using these techniques the study material will provide a unique record to study (changes in) rainfall dynamics of up to 900 years in this region strongly susceptible for climate change. Integration with already existing networks of juniper chronologies in Pakistan (Esper 2000, Treydte et al. 2006) and Tibet (Bräuning 1999, 2001) and other (paleo)records, such as speleothems (Fleitmann et al. 2002, 2003, 2007) and varves and turbidites (Berger & Rad 2002) will contribute to a better understanding of large scale climate circulation patterns and their changes on the Arabian Peninsula throughout the last millennium. For Oman a reconstruction of rainfall will be of paramount importance to further elucidate the historic development of oasis agriculture.

### **Acknowledgements**

The authors are indebted to Maher Nagieb for collecting many of the tree cores and to the Deutsche Forschungsgemeinschaft for funding (BU 1308).

### **References**

- Berger, W.H., Rad, U., von (2002): Decadal to millennial cyclicity in varves and turbidites from the Arabian sea: hypothesis of tidal origin. *Global and Planetary Change* 34: 313-325.
- Bräuning, A. (1999): Dendroclimatological potential of drought-sensitive tree stands in Southern Tibet for the reconstruction of monsoonal activity. *IAWA Journal* 20(3): 325-338.

- Bräuning, A. (2001): Climate history of the Tibetan Plateau during the last 1000 years derived from a network of juniper chronologies. *Dendrochronologia* 19:127-37.
- Cherubini, P., Gartner, B.L., Tognetti, R., Bräker, O.U., Schoch, W., Innes, J.L. (2003): Identification, measurement and interpretation of tree rings in woody species from mediterranean climates. *Biol. Rev.* 78: 119-148.
- Cook, E.R., Kairiukstis, L.A. (1990): Methods of Dendrochronology: Applications in the Environmental Sciences. International Institute for Applied Systems Analysis, *Kluwer Academic Publishers*, Boston, MA, USA.
- Courealet, C., Sass-Klaassen, U., Sterck, F., Bekele, T., Zuidema, P.A. (2005): Combining dendrochronology and matrix modelling in demographic studies: An evaluation for *Juniperus procera* in Ethiopia. *Forest Ecology and Management* 216: 317-330.
- Esper, J. (2000): Long-term tree-ring variations in *Juniperus* at the upper timber-line in Karakorum (Pakistan). *The Holocene* 10: 253-60.
- Fisher, M. (1994): Is it possible to construct a tree-ring chronology for *Juniperus excelsa* (Bieb.) subsp. *polycarpus* (K. Koch), Takhtajan from the northern mountains of Oman? *Dendrochronologia* 12: 119-127.
- Fisher, M., Gardner, A.S. (1995): The status and ecology of a *Juniperus excelsa* subsp. *polycarpus* woodland in the northern mountains of Oman. *Vegetation* 119: 33-51.
- Fichtler, E., Trouet, V., Beeckman, H., Coppin, P., Worbes, M. (2004): Climatic signals in tree rings of *Burkea africana* and *Pterocarpus angolensis* from semiarid forests in Namibia. *Trees* 18, 442–51.
- Fleitmann, D., Burns, S.J., Neff, U., Al-Subbary, A.A., Al-Aowah, M.A., Matter, A., Mangini, A. (2002): Climate change in Southern Arabia: Speleothems from Yemen and Oman as archives for monsoon rainfall over the last 330,000 years. *Yemeni Journal of Science* 4: 15-23.
- Fleitmann, D., Burns, S.J., Mudelsee, M., Neff, U., Kramers, J., Mangini, A., Matter, A. (2003): Holocene forcing of the Indian monsoon recorded in a stalagmite from Southern Oman. *Science* 300: 1737-1739.
- Fleitmann, D., Burns, S.J., Mangini, A., Mudelsee, M., Kramers, J., Villa, I., Neff, U., Al-Subbary, A.A., Buettner, A., Hippler, D., Matter, A., 2007. Holocene ITCZ and Indian monsoon dynamics recorded in stalagmites from Oman and Yemen (Socotra). *Quaternary Science Reviews* 26, 170–188.
- Kuzmin, Y.V., Slusarenko, I.Y., Hajdas, I. Bonani, G. Christen, A. (2004): The Comparison of 14C Wiggle-Matching Results for the 'Floating' Tree-Ring Chronology of the *Ulandryk-4 Burial Ground (Altai Mountains, Siberia)*. *Radiocarbon* 46: 943-948.
- von Rad, U., Schaaf, M., Michels, K.H., Schulz, H., Berger, W.H., Sirocko, F. (1999): A 5000-yr Record of Climate Change in Varved Sediments from the Oxygen Minimum Zone off Pakistan, Northeastern Arabian Sea. *Quaternary Research* 51 (1): 39-53.
- Schöngart, J.B., Orthmann, K.J., Hennenberg, S., Porembski, Worbes, S. (2006): Climate-growth relationships of tropical tree species in West Africa and their potential for climate reconstruction. *Global Change Biology* 12: 1139-50.

- Siebert, S., Häser, J., Nagieb, M., Korn, L., Buerkert, A. (2005): Agricultural, architectural and archaeological evidence for the role and ecological adaptation of a scattered mountain oasis in Oman. *Journal of Arid Environments* 62: 177-197.
- Trouet, V., Coppin, P., Beeckman, H. (2006): Annual growth ring patterns in *Brachystegia spiciformis* reveal influence of precipitation on tree growth. *Biotropica* 38, 375-82.
- Treydte, K.S., Schleser, G.H., Helle, G., Frank, D.C., Winiger, M., Haug, G.H., Esper, J. (2006): The twentieth century was the wettest period in northern Pakistan over the past millennium. *Nature* 440 (27), 1179-1182.
- Wils, T.H.G., Eshetu, Z. (2007): Reconstructing the flow of the River Nile from *Juniperus procera* and *Prunus africana* tree rings (Ethiopia) – an explorative study on cross-dating and climate signal. *TRACE, Vol. 5*, 277-284.
- Worbes, M. (1995): How to measure growth dynamics in tropical trees, a review. *IAWA Journal* 16 (4): 337-351.